


Please type a plus (+) inside this box - 

PTO/SB/29 (12/97)

Approved for use through 09/30/00. OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

# UTILITY PATENT APPLICATION TRANSMITTAL

Attorney Docket No. 9114-004-999 Total Pages 35

First Named Inventor or Application Identifier

Dennis Mangano

Express Mail Label No. EL 452 479 835 US

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

## APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner for Patents  
Box Patent Application  
Washington, DC 20231

1. ☒ Fee Transmittal Form  
*Submit an original, and a duplicate for fee processing*
2. ☒ Specification [Total Pages 28]  
*(preferred arrangement set forth below)*
  - Descriptive title of the Invention
  - Cross Reference to Related Applications
  - Statement Regarding Fed sponsored R&D
  - Reference to Microfiche Appendix
  - Background of the Invention
  - Brief Summary of the Invention
  - Brief Description of the Drawings *(if filed)*
  - Claim(s)
  - Abstract of the Disclosure
3. ☒ Drawing(s) (35 USC 113) [Total Sheets 2]
4. ☒ Oath or Declaration [Total Sheets 2]
  - a. ☐ Newly executed (original or copy)
  - b. ☒ Copy from a prior application (37 CFR 1.63(d))  
*(for continuation/divisional with Box 17 completed)*  
*[Note Box 5 below]*
    - i. ☐ DELETION OF INVENTOR(S)  
Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33 (b).
5. ☒ Incorporation By Reference *(useable if Box 4b is checked)*
6. ☐ Microfiche Computer Program *(Appendix)*
7. ☐ Nucleotide and/or Amino Acid Sequence Submission *(if applicable, all necessary)*
  - a. ☐ Computer Readable Copy
  - b. ☐ Paper Copy (identical to computer copy)
  - c. ☐ Statement verifying identity of above copies

## ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & document(s))
9. ☐ 37 CFR 3.73(b) Statement ☐ Power of Attorney  
*(when there is an assignee)*
10. ☐ English Translation Document *(if applicable)*
11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard (MPEP 503)  
*(Should be specifically itemized)*
14. ☐ Small Entity ☐ Statement filed in prior application, Statement(s) Status still proper and desired
15. ☐ Certified Copy of Priority Document(s)  
*(if foreign priority is claimed)*
16. ☐ Other:

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☒ Continuation ☐ Divisional ☐ Continuation-in-part (CIP)

of prior application no: 08/787,056 filed December 3, 1996.

## 18. CORRESPONDENCE ADDRESS

☒ Customer Number or Bar Code Label

20583

*(Insert Customer No. or Attach bar code label here)*

or ☐ Correspondence address below

NAME					
ADDRESS					
CITY	STATE	ZIP CODE			
COUNTRY	TELEPHONE	FAX			

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

**PENNIE & EDMONDS LLP**

COUNSELLORS AT LAW

1155 Avenue of the Americas

New York, N.Y. 10036-2711

(212) 790-9090

ATTORNEY DOCKET NO. 9114-004-999Date: October 22, 1999

Assistant Commissioner for Patents  
 Box PATENT APPLICATION  
 Washington, D.C. 20231

Sir:

The following utility patent application is enclosed for filing:

Applicant(s): Dennis Mangano

Executed on: June 14, 1997

Title of Invention: METHODS FOR REDUCING MORTALITY AN MORBIDITY BY POSTOPERATIVE  
 ADMINISTRATION OF A PHARMACOLOGIC CARDIOVASCULAR AGENT

**PATENT APPLICATION FEE VALUE**

TYPE	NO. FILED	LESS	EXTRA	EXTRA RATE	FEE
Total Claims	48	-20	28	\$18.00 each	\$ 504.00
Independent	3	-3	0	\$78.00 each	\$ 0.00
Minimum Fee					\$ 760.00
Multiple Dependency Fee If Applicable (\$260.00)					\$ 0.00
<b>Total</b>					\$ 1,264.00
50% Reduction for Independent Inventor, Nonprofit Organization or Small Business Concern (a verified statement as to the applicant's status is attached)					- \$ 632.00
<b>Total Filing Fee</b>					\$ 632.00

- ☒ Priority of application no. 08/787,056 filed on December 3, 1996 in is claimed under 35 U.S.C. § 119.  
☐ The certified copy of the priority application has been filed in application no. filed  
☐ Amend the specification by inserting before the first line the following sentence: This is a continuation-in-part of application no. filed .

Please charge the required fee to Pennie & Edmonds LLP Deposit Account No. 16-1150. A copy of this sheet is enclosed.

Respectfully submitted,

*Samuel B. Abrams* 40,611  
 Samuel B. Abrams 30,605  
 PENNIE & EDMONDS LLP (Reg. No.)

Enclosure

This form is not for use with continuation, divisional, re-issue, design or plant patent applications.

**METHODS FOR REDUCING MORTALITY  
AND MORBIDITY BY POSTOPERATIVE ADMINISTRATION  
OF A PHARMACOLOGIC CARDIOVASCULAR AGENT**

**1. INTRODUCTION**

5       The present invention relates to methods for reducing mortality and cardiovascular morbidity following surgery. In particular, the invention relates to the intensive postoperative administration of a pharmacologic cardiovascular agent to reduce mortality and cardiovascular complications. The invention is illustrated by way of  
10 working examples which demonstrate that in patients with, or at risk for, coronary artery disease undergoing major noncardiac surgery, the administration of a  $\beta$ -adrenergic blocking agent throughout the period of hospitalization:  
1) reduces mortality and cardiovascular complications following hospital discharge; 2) is safe and well tolerated;  
15 and 3) the estimated cost savings in lives more than outweighs the cost of therapy.

**2. BACKGROUND OF THE INVENTION**

Cardiovascular mortality and morbidity are prevalent and costly for the 30 million patients undergoing noncardiac  
20 surgery annually in the United States. More than one million of these patients suffer heart attacks or other cardiac complications after the operation, with about 500,000 resultant deaths during the first two postoperative years (Mangano, 1990, Anesthesiology 72: 153-184; Mangano and Goldman, 1995, N. Eng. J. Med. 333:1750-1756). In the subset  
25 of 3 million surgical patients with or at-risk for coronary artery disease, the most significant risk factors for mortality and cardiovascular morbidity are myocardial ischemia and non-fatal myocardial infarction occurring during the first week following surgery, which increases the risk of serious adverse cardiovascular outcomes by 2- to 20-fold over  
30 the two years following surgery (Mangano et al., 1990, N. Eng. J. Med. 323: 1781-1788; Mangano et al., 1992, JAMA

268:233-239; Browner et al., 1992, JAMA 268:228-232). These postoperative ischemic events appear to be related to the persistent exaggerated sympathetic response that is associated with substantial increases in heart rate throughout the in-hospital period (Rao et al, 1983

- 5 Anesthesiology 59:499-505; Gottlieb et al., 1987, J. Am Coll. Cardiol. 10:756-760; Siliciano et al., 1990, Postoperative Myocardial Infarction: Mechanisms and Therapies, In Estafanous (ed.): Opioids in Anesthesia, Butterworth Publishers, Boston pp. 164-177; Mangano et al., 1991, J. Am Coll. Cardiol. 17:843-850; Mangano et al., 1991, J. Am. Coll. Cardiol. 17:851-857).

Studies conducted over the past decade have established the association between postoperative myocardial ischemia and post-discharge adverse outcomes, with the odds of such outcomes increasing in patients with (versus without) postoperative ischemia by 28-fold six months following

- 15 surgery, 20-fold at one year, and 14-fold at two years (Mangano et al., 1990, N. Eng. J. Med. 323:1781-1788; Mangano et al., 1992, JAMA 268:233-239; Browner et al., 1992, JAMA 268:228-232; Raby et al., 1989, N. Eng. J. Med. 321:1296-1300; Slogoff and Keats, 1985, Anesthesiology 62:107-114; Eisenberg et al., 1992, JAMA 268:210-216). In addition,
- 20 studies have demonstrated an association between postoperative ischemia and elevated heart rate, and have suggested that mitigation of this heart rate response may reduce the incidence and/or severity of ischemia (Rao et al. 1983, Anesthesiology 59:499-505; Gottlieb et al., 1987, J. Am. Coll. Cardiol. 10:756-760; Mangano et al., 1991, J. Am. Coll. Cardiol. 17:843-850; Mangano et al., 1991, J. Am Coll. Cardiol. 17:851-857; Wallace et al., 1994, Anesthesiology 81:A99).

- In at-risk patients about to undergo major surgery, the standard practice is to control heart rate prior to surgery, to continue medication to the time of surgery, and to
- 30 modulate the heart rate response during surgery using anesthetic techniques. However, following surgery heart rate

is not well-controlled, increasing above preoperative levels by 30 percent or more, throughout the extended postoperative period (Mangano et al., 1992, JAMA 268:233-239; Mangano et al., 1991, J. Am. Coll. Cardiol. 17:843-850; Mangano et al., 1991, J. Am. Coll. Cardiol. 17:851-857; Raby et al., 1989, N. Eng. J. Med. 321:1296-1300; Eisenberg et al., 1992, JAMA 268:210-216). Furthermore, even brief periods of tachycardia during the postoperative period may precipitate ischemia in these patients, who also are subjected to alterations in perfusion, oxygenation and coagulation, as well as other stresses imposed by the exaggerated sympathetic and inflammatory responses to surgery. However, despite appreciation of the general problem of perioperative infarction, as well as the potentially deleterious effect of an unchecked postoperative sympathetic response, and despite recognition of the efficacy of  $\beta$ -blockade in ambulatory patients with coronary artery disease, clinicians have been reluctant to prescribe  $\beta$ -blockers following surgery, even for patients who had been maintained on  $\beta$ -blockers prior to admission for surgery. Such reluctance is based on several concerns, including: 1) safety - namely precipitation of postoperative heart failure, hypotension and bronchospasm; 2) efficacy - unproven for the surgical patient; and 3) cost.

Several clinical trials have investigated the effects of preoperative or intraoperative use of nitrates (Coriat et al., 1984, Anesthesiology 63: 193-196; Gallagher et al., 1986, Anesthesiology 64:785-789),  $\beta$ -adrenergic blockers (Stone et al., 1988, Anesthesiology 68:495-500; Magnusson et al., 1986, Br. J. Anaeth. 58:251-260; Cucchiara et al., 1986, 65-528-531), calcium channel blockers (Chung et al., 1988, Anesthesiology 69:343-347; Merin, 1987, Anesthesiology 66:111), and alpha ( $\alpha$ )-2 agonists (Ghignone et al., 1987, Anesthesiology 67:3-10; Talke et al., 1995, Anesthesiology 82:629-633) on hemodynamics and measures of myocardial ischemia. In the studies involving  $\beta$ -blockers, the drugs were always administered prior to the surgical procedures. Prior to the present invention, it was not known that

continuous postoperative administration of these agents would result in a reduction of cardiovascular mortality and morbidity. In particular, it was not expected that it would have any long-term beneficial effects on mortality and cardiovascular events, such as myocardial infarction, heart failure and unstable angina requiring revascularization.

### 3. SUMMARY OF THE INVENTION

The present invention relates to methods for reducing mortality and cardiovascular morbidity following surgery by the intraoperative and postoperative administration of a therapeutic amount of a pharmacologic cardiovascular agent. In particular, it relates to the intensive postoperative administration of such an agent during hospitalization and even after hospital discharge to mitigate the sympathetic response associated with increased heart rate, increased thrombosis and increased inflammatory response, thereby reducing the incidence and/or severity of cardiovascular complications such as myocardial infarction, unstable angina, congestive heart failure, dysrhythmia, myocardial revascularization, and death.

The invention is based, in part, on the Applicant's discovery that the administration of a  $\beta$ -adrenergic blocker, atenolol, prior to and immediately following surgery, and continuing daily throughout the entire period of hospitalization in patients with, or at risk for, coronary artery disease undergoing noncardiac surgery, reduces mortality and serious cardiovascular complications following hospital discharge, with the early survival effects persisting for two years. Therefore, a wide variety of uses are encompassed by the present invention including, but not limited to, increasing the survival rate and decreasing cardiovascular complications in patients under surgical stress.

4. BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Intensive postoperative administration of atenolol increases the survival of patients for two years after surgery.

5 Figure 2. Intensive postoperative administration of atenolol increases cardiovascular event-free survival of patients for two years after surgery.

5. DETAILED DESCRIPTION OF THE INVENTION

10 The present invention relates to treatment of patients undergoing surgery to reduce mortality and cardiovascular complications by the administration of a pharmacologic cardiovascular agent such as a  $\beta$ -adrenergic blocking agent following surgery. The treatment may be continued throughout hospitalization, and even after discharge. Both the time-to-  
15 first adverse event, as well as survival and event-free survival, are significantly improved by such treatment, particularly during the first 6-8 months following surgery, with survival effects persisting for two years. In the  $\beta$ -adrenergic blocking agent-treated patients, survival was 90 percent at two years following surgery versus 79 percent in  
20 placebo-treated patients, and event-free survival was 83 percent versus 68 percent, respectively. Moreover, the intensive postoperative drug administration was well-tolerated in these patients, despite the prevalence of cardiac and pulmonary disease.

25 The invention is discussed in more detail in the subsections below, solely for the purpose of description, and not by way of limitation. Although the specific procedures and methods described herein are exemplified with the administration of atenolol immediately before and after surgery and continuing for up to seven days thereafter, they are merely illustrative for the practice of the invention.  
30 Analogous schedules, procedures, techniques and pharmacologic cardiovascular agents are equally applicable.

### 5.1 SUITABLE PHARMACOLOGIC CARDIOVASCULAR AGENTS

The present invention relates to the intensive postoperative use of a pharmacologic cardiovascular agent to reduce mortality and morbidity following surgery. As used herein, a "pharmacologic cardiovascular agent" is an agent that mitigates cardiovascular stress responses by reducing heart rate, blood coagulation or inflammatory reactions. In a specific embodiment of the invention by way of working examples, *infra*, a  $\beta$ -adrenergic blocking agent is used to reduce heart rate.  $\beta$ -adrenergic receptors are expressed on different cell types, including cardiac muscle cells. These receptors are further subdivided into  $\beta_1$  and  $\beta_2$  receptors on the basis of their tissue distribution, both forms are coupled to a signal transducer referred to as the G protein. The binding of these receptors by a ligand results in G-protein-mediated activation of the enzyme adenylate cyclase, which causes an elevation of intracellular cyclic AMP levels as well as activities of ion channels. An increase of cyclic AMP regulates a number of downstream cellular metabolic events. Such events are manifested in an increased contraction rate of cardiac muscle cells which, in turn, promotes increased heart rate and blood pressure. Under certain circumstances of bodily stress such as surgery, these events can lead to serious cardiovascular complications, even death. In view of the foregoing observation, a number of  $\beta$ -adrenergic blocking agents or antagonists have been tested clinically for the treatment of hypertension, ischemic heart disease and certain cardiac arrhythmias.

The interactions between hormones such as epinephrine and  $\beta$ -adrenergic receptors have been well studied in the art. The binding of epinephrine to  $\beta$ -adrenergic receptors activates adenylate cyclase, and a number of measurable downstream cellular events. Since the  $\beta$ -adrenergic receptor has been molecularly cloned and expressed in receptor-negative cells, the ability of such cells to activate adenylate cyclase in response to epinephrine may be



conveniently tested in an *in vitro* assay system.

Alternatively, a  $\beta$ -adrenergic receptor-positive cell line may also be used. For the purpose of this invention, any substance that blocks or interferes with the activation of adenylate cyclase by a ligand such as epinephrine upon its binding to the  $\beta$ -adrenergic receptors is a  $\beta$ -adrenergic blocking agent suitable for use in the present invention.

In accordance with the methods of the invention,  $\beta$ -adrenergic blocking agents encompass both  $\beta_1$ -selective and non-selective blockers. However,  $\beta_1$ -selective blockers are preferred because they exert minimal effects on the  $\beta$ -adrenergic receptors on non-cardiac muscle cells. Examples of suitable blocking agents include, but are not limited to, atenolol, metoprolol, esmolol, acebutolol, practolol, alprenolol, propranolol, nadolol, timolol, pindolol, labetalol, sotalol and oxprenolol. The aforementioned agents are commercially available or may be readily prepared by methods well known in the art (Goodman and Gilman's, The Pharmacological Basis of Therapeutics, 1990, eighth ed., Pergamon Press).

Additionally, other pharmacologic agents with known cardiovascular effects in reducing heart rate, blood coagulation and inflammation are also suitable for use in the present invention, and such agents include, but are not limited to,  $\alpha$ -2 agonists such as clonidine, anti-ischemic agents which encompass calcium channel blockers such as verapamil and nifedipine, angiotensin converting enzyme (ACE) inhibitors such as lisinopril and enalapril, and nitrates (nitroglycerin), antiplatelet agents such as aspirin and dipyridamole, antithrombotics such as coumadin, heparin and streptokinase, and the like (Physicians' Desk Reference, 1996, 50th Edition, Medical Economics).

## **5.2 DOSAGE AND FORMULATION**

The agents described in Section 5.1, *supra*, may be administered into a patient for the reduction of mortality

and cardiovascular morbidity following surgery by any means that produces contact of the active agent with the agent's site of action in the body of the patient. They can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual  
5 therapeutic agents or in a combination of therapeutic agents. Each can be administered alone but is generally administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice. The pharmaceutical compositions of the invention may be adapted for oral, parenteral or topical  
10 administration, and may be in unit dosage form, in a manner well known to those skilled in the pharmaceutical art. Parenteral administration includes, but is not limited to, injection subcutaneously, intravenously, intraperitoneally or intramuscularly.

In accordance with this aspect of the invention, a  
15 pharmacologic cardiovascular agent is administered during and after the surgical operation, and if tolerated by the patient, it may be continued for 3-7 days until the patient is discharged from the hospital, and following hospital discharge. The administration may begin immediately after surgery and continuing daily through hospital discharge and  
20 following hospital discharge. Alternatively, the administration may be initiated only after the first clinical manifestation of cardiovascular stress such as high blood pressure, hypertension, myocardial ischemia or infarction, increased heart rate relative to the preoperative rate, clotting abnormalities and inflammatory reaction such as a  
25 rise in body temperature, or a local tissue reaction involving infiltration of white blood cells or other inflammatory mediators. In addition, the administration may begin prior to surgery, with the preferred timing of administration being from 1 day to 1 hour before surgery.

The dose administered will, of course, vary depending  
30 upon known factors, such as: the pharmacodynamic characteristics of the particular agent and its mode and

route of administration; the age, health, height and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment(s); the frequency of treatment(s); and the effect desired. A daily dose of active ingredient can be expected to be about 0.1-500 mg per

5 patient, with the preferred dose being 5-100 mg given in two separate doses.

More specifically, atenolol may be used orally at 50-100 mg/day and at 5-10 mg BID intravenously. Labetalol may be used orally at 200-400 mg BID and at 20 mg bolus intravenously over 2 minutes with a repeated dose of 40-80 mg  
10 over 10 minutes up to a maximum dose of 300 mg. Clonidine may be used orally at 0.2-1.2 mg/day and at 0.1-0.3 mg by skin patch every 7 days. Nitroglycerine may be used orally at gr 1/100-gr 1/400 sublingual with a repeated dose for 15-30 minutes and at 1-10µg/kg/minute. Verapamil may be used orally at 240-320 mg/day and at 0.075-0.15 mg/kg  
15 intravenously over 2 minutes. Nifedipine may be used orally at 10-20 mg TID. Lisinopril may be used orally at 20-40 mg/day. Enalapril may be used orally at 10-40 mg/day and at 1.25 mg intravenously for 6 hours. Aspirin may be used orally at 325-650 mg/day or BID. Dipyridamole may be used orally at 50-400 mg/day and at 0.142 mg/kg/minute  
20 intravenously over 4 minutes up to a total dose of 0.5 mg/kg. Coumadin may be used orally at 10-15 mg/day for 3 days followed by 2-3 mg/day. Heparin may be used at 1000-5000U intravenous push or 5000-7500U intravenous bolus followed by adjustment according to PTT. Streptokinase may be used intravenously at 20,000 IU bolus followed by a dose of 2,000  
25 IU/minute for 60 minutes.

Dosage forms (compositions suitable for administration) contain from about 1 mg to about 500 mg of active ingredient per unit. In these pharmaceutical compositions, the active ingredient is ordinarily present in an amount of about 0.5-95% by weight based on the total weight of the  
30 composition.

The active ingredient can be administered orally in solid or semi-solid dosage forms, such as hard or soft-gelatin capsules, tablets, or powders, or in liquid dosage forms, such as elixirs, syrups, or suspensions. It can also be administered parenterally, in sterile liquid dosage forms.

- 5 Other dosage forms are potentially possible such as patches or ointment or transdermal administration.

Gelatin capsules or liquid-filled soft gelatin capsules may contain the active ingredient and powdered or liquid carriers, such as lactose, lecithin starch, cellulose derivatives, magnesium stearate, stearic acid, and the like.

- 10 Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar-coated or film-coated to mask any unpleasant taste and to protect the tablet from the atmosphere, or enteric-coated  
15 for selective disintegration in the gastrointestinal tract. Liquid dosage forms for oral administration can contain coloring and/or flavoring to increase patient acceptance.

- In general, water, oil, saline, aqueous dextrose (glucose), polysorbate and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are  
20 suitable carriers for parenteral solutions. Solutions or emulsions for parenteral administration preferably contain about 5-15% polysorbate 80 or lecithin, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents including, but not limited to, sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined,  
25 are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, including but not limited to, benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol.

- Suitable pharmaceutical carriers are further described  
30 in *Remington's Pharmaceutical Sciences*, 1990, 17th ed., Mack

Publishing Company, Easton, PA, a standard reference text in this field, which is incorporated herein by reference in its entirety.

**5.3 METHODS OF USING PHARMACOLOGIC CARDIOVASCULAR  
AGENTS IN PATIENTS UNDERGOING SURGERY**

5  
10 The methods of the invention are generally applicable to patients undergoing surgery to reduce their long-term mortality and cardiovascular morbidity. The methods are particularly useful for patients who have or are at risk for coronary artery disease. The cardiac risk factors include hypertension, smoking, diabetes mellitus, age over 65 and cholesterol level  $>6.2$  mmol/liter.

15 The methods of the invention are applicable to patients undergoing any form of surgery that causes cardiovascular stress whether or not it is cardiac surgery. The types of surgery include, but are not limited to, intraabdominal, orthopedic, neurological, intrathoracic, head and neck, vascular and general surgery.

20 Intensive administration of the pharmacologic cardiovascular agents may begin immediately after surgery. However, in the case of coronary bypass, it may begin intraoperatively, either prior to or following institution of cardio-pulmonary bypass or at any time during postoperative hospitalization. Alternatively, initiation of therapy may await the first manifestations of cardiovascular stress. Markers for such stress include high blood pressure, hypertension, myocardial infarction, unstable angina, tachycardia, clotting abnormalities and inflammatory  
25 response. Intensive therapy refers to daily administration of the pharmacologic agents until reduction of symptoms of cardiovascular stress or hospital discharge.

30

6. **EXAMPLE: ADMINISTRATION OF A BETA-BLOCKER REDUCES MORTALITY AND CARDIOVASCULAR MORBIDITY FOLLOWING SURGERY**

6.1 **MATERIALS AND METHODS**

6.1.1 **PATIENT POPULATION**

5 Eligible patients included those with, or at risk for coronary artery disease and scheduled for elective noncardiac surgery requiring general anesthesia at the San Francisco Veterans Affairs Medical Center. The specific inclusion and exclusion criteria have been described previously (Mangano et al., 1990, N. Eng. J. Med. 323:1781-10 1788; Mangano et al., 1992, JAMA 268:233-239). A maximum of one patient per day was enrolled and, of the 204 patients consenting to the study, 200 were enrolled randomized and received study drug.

6.1.2 **BETA-ADRENERGIC BLOCKING AGENT**

15 Patients were randomized to receive either atenolol "TENORMIN" (Zeneca Pharmaceuticals) or placebo prior to induction of anesthesia, immediately following surgery, and daily throughout their hospitalization (up to 7 days). Drug assignment, study physicians, treating clinicians, and data analysis personnel were blinded to study group throughout all 20 phases of this trial. Intravenous and oral preparations of active drug atenolol and placebo were prepared by the hospital pharmacy with a computer-generated randomized list retained only by the pharmacy and maintained confidential until formal study unblinding following database closure.

25 Intravenous preparation consisted of two-10-ml syringes, each containing 5 mg atenolol or placebo; oral preparation consisted of two 50 mg tablets of atenolol, or two placebo tablets. Approximately one hour prior to surgery, patients entered the preoperative area and blood pressure was recorded with an automated cuff and 5-lead 30 continuous electrocardiograph. Thirty-minutes prior to entry into the operating room, intravenous administration of study

drug began. Exclusion criteria for study drug administration were heart rate <55 bpm, systolic blood pressure <100 mm Hg, or evidence of congestive heart failure, third degree heart block, or bronchospasm (ISIS-I protocol, 1986, Lancet 2:56-66). If none of these criteria was present, the first  
5 syringe of study drug was infused over five minutes, the patient was observed for an additional five minutes, and, if no exclusion criteria developed, the second syringe was infused over five minutes. Immediately following surgery, the study drug was again given using the identical technique applied prior to surgery. On the morning of the first  
10 postoperative day, and daily thereafter until the patient was discharged from the hospital (up to a maximum of seven days), patients received study drug every 12 hours using the same technique for intravenous infusion, or orally (if able) at which time a tablet of atenolol (0, 50 or 100 mg) or placebo was given daily. If heart rate was >65 bpm and systolic  
15 blood pressure >100 mm Hg, 100 mg atenolol (or placebo) was given orally; if heart rate was >55 but ≤65 bpm and systolic blood pressure >100 mm Hg, 50 mg atenolol (or placebo) was administered; if heart rate was <55 bpm or systolic blood pressure <100 mm Hg, 0 mg atenolol (or placebo) was given. No treating clinician was allowed to observe study drug  
20 administration either prior to, or after, surgery.

### 6.1.3 CLINICAL CARE

All patients received general anesthesia with endotracheal intubation; preoperative medications were continued until the time of surgery, with beta-blockers  
25 replaced by study drug on the morning of surgery. There were no other protocol-based restrictions of anesthetic or surgical technique, and clinical decisions were not controlled by study protocol. Perioperative information was recorded and analyzed for possible confounding effects, and included: type and duration of surgery, anesthetic  
30 techniques, fluid and blood loss and replacement,

cardiovascular medications, hemodynamics,  
electrocardiographic data, and adverse events.

#### 6.1.4 CLINICAL FOLLOW UP AND OUTCOME MEASUREMENTS

5 Of the 200 patients enrolled, 194 were discharged  
following surgery and six patients died during hospitalization  
- three cardiac deaths secondary to myocardial infarction  
(two placebo and one atenolol), and three noncardiac deaths,  
with two secondary to metastatic cancer (both atenolol), and  
10 one with pulmonary failure secondary to massive infusion for  
fluid loss (atenolol). Of the 194 patients discharged,  
outcome data were collected in 192 patients (99%), with two  
patients (one placebo and one atenolol) lost to followup. At  
six months, one year and two years following surgery, study  
physicians conducted scheduled research study visits that  
15 were independent of usual clinical care. Data collected  
during each visit included history and physical examination,  
a 12-lead electrocardiogram, and review of all medical  
records, medications and hospital admissions. Cardiac death  
was diagnosed if the patient died of either a myocardial  
infarction, dysrhythmia or congestive heart failure caused  
primarily by a cardiac condition. Myocardial infarction  
20 required the following: 1) development of new Q waves (as  
defined by Minnesota Code 1-1-1-21-2-7); or 2) new persistent  
ST-T wave changes (Minnesota Code 4-1 or 4-2; 5-1 or 5-2)  
associated at the time of hospitalization with elevation of  
total creatinine kinase and CK-MB isoenzyme; or 3) necropsy  
evidence of acute myocardial infarction; or 4) hospital  
25 record documentation of myocardial infarction (Mangano et  
al., 1990, N. Eng. J. Med. 323:1781-1788). Unstable angina  
required severe precordial chest pain that lasted at least 30  
minutes, was unresponsive to standard therapeutic maneuvers  
and associated with transient ST-segment or T-wave changes  
without development of Q waves or diagnostic enzyme  
30 abnormalities. The diagnosis of congestive heart failure



required symptoms or signs of pulmonary congestion (shortness of breath and rales), signs of new left or right ventricular failure (cardiomegaly, S3, jugular venous distention, and peripheral edema), abnormal results on chest radiography (vascular redistribution, interstitial edema, and alveolar edema), and a change in medication involving (at least) treatment with diuretic agents (Mangano et al., 1990, N. Eng. J. Med. 323:1781-1788).

Outcomes were prescribed by study protocol, and the primary outcome was all-cause mortality during the two years following hospital discharge. The secondary outcome was combined consisting of : 1) myocardial infarction; or 2) unstable angina or congestive heart failure requiring hospital admission and clinical diagnosis and treatment, or 3) myocardial revascularization (coronary artery bypass graft surgery or percutaneous transluminal angioplasty), or 4) death. Autopsy data, if available for patients who died over the two-year period, were reviewed centrally at the core laboratory (Ischemia Research and Education Foundation) by a pathologist blinded to patient treatment group.

#### 6.1.5 STATISTICAL ANALYSIS

The study was designed to allow assessment of tolerance, in-hospital events (hemodynamic changes, dysrhythmia, ischemia), and adverse cardiovascular outcomes occurring over the two years following surgery. A sample size of 200 patients was calculated based on the following assumptions: 1) duration of enrollment and followup = 48 months; 2) two-year mortality, cardiovascular morbidity and in-hospital event rates = 0.23, 0.28 and 0.41, respectively (Mangano et al., 1990, N. Eng. J. Med. 323:1781-1788; Mangano et al., 1992, JAMA 268:233-239; Browner et al., 1992, JAMA 268:228-232); 3) followup rate = 0.96 (Mangano et al., 1992, JAMA 268:233-239; Browner et al. 1992, JAMA 268:228-232); 4)  $\alpha=0.05$ ,  $\beta=0.2$ , effect size = 0.5; and 5) the alternative safety and efficacy hypotheses are two-tailed and one-tailed,

respectively (Mangano, 1990, Anesthesiology 72:153-184; Stone et al., 1988, Anesthesiology 68:495-500; Magnusson et al., 1986, Br. J. Anaesth. 58:251-260; Cucchiara et al., 1986, Anesthesiology 65:528-531; Wallace et al., 1994, Anesthesiology 81:A99). Using the log-rank survival test for  
5 sample size estimation (BMDP statistical Software Inc., 1992, SOLO Power Analysis), it was calculated that 198 patients would be necessary for mortality assessment and 158 patients for combined outcome, and, using Z statistic, 170 patients for in-hospital event assessments. Mortality risk in  
10 different categories (all-cause mortality, cardiac mortality, noncardiac mortality, at 6 months, 1 year, and 2 years) was compared using Kaplan-Meier methods, as was event-free survival after discharge. Univariable predictors of two-year mortality were identified using the Cox proportional hazards regression techniques (SAS Institute Inc., 1992, Release 6.07:345-379) after first verifying that assumption of the  
15 hazards model was valid (Kaplan and Meier, 1958, J. Am. Stat. Assoc. 53: 457-481; SAS Institute, Inc., 1985, Statistical Analysis System, SAS User's Guide). Predictors with  $P < 0.10$  were entered into multivariable models and a series of models was constructed by adding variables, as long as the resulting multivariable model had a lower Chi-square P value than  
20 competing models. Analyses were performed using Statistical Analysis System Software (SAS Institute, Inc., Cary, NC).

## 6.2 RESULTS

Study patients were middle-aged or elderly who smoked and had a history of hypertension and chronic medical  
25 problems. There was no difference between groups, except that the atenolol group had a higher incidence of treatment for hypertension.

Thirty patients (15.6 percent) died over the two-year outcome period (Table 1). Twenty-one of these deaths (12 cardiac-related) occurred in the placebo group versus 9  
30 deaths (4 cardiac-related) in the atenolol group,

representing a 57 percent reduction by atenolol in all-cause mortality ( $P=0.019$ ), and a 67 percent reduction in cardiac mortality ( $P=0.033$ ). The principal effect of atenolol therapy was on cardiac-related outcomes occurring over the first 6-8 months (one noncardiac death versus 10 deaths with 5 7 cardiac-related;  $P<0.001$ ), with the time to first death being 19 days in the placebo group versus 237 days in the atenolol group. Thereafter there was no substantial effect; however, the early difference in survival between groups was preserved at one year (3 versus 14 deaths;  $P = 0.005$ ) and two years (9 versus 21 deaths;  $P = 0.019$ ), with survival 10 significantly increased over all time periods in the atenolol group (Figure 1).

Atenolol-treated patients had a significant decrease in the rate of cardiac events within six months following surgery (0 atenolol patients versus 12 placebo patients;  $P < 0.001$ ), a 3-fold decrease within one year (7 atenolol 15 patients versus 22 placebo patients;  $P = 0.003$ ), and a 2-fold decrease within two years following surgery (16 atenolol patients versus 32 placebo patients;  $P = 0.008$ ). The principal effect occurred over the first 6 to 8 months, with the time-to-first adverse event being 6 days in the placebo group versus 158 days in the atenolol group. Thereafter, 20 there was no substantial effect; however, the early difference in event-free survival was preserved over the two years following surgery (Figure 2).

During treatment, the average heart rate was significantly lower in the atenolol group (75 bpm versus 87 bpm;  $P < 0.001$ ), as was the maximum heart rate (113 bpm versus 25 130 bpm;  $P < 0.001$ ). Multivariable correlates associated with survival at two years are listed in Table 2, and demonstrate association between survival and a history of diabetes mellitus and atenolol therapy, with atenolol improving survival in diabetics at two years by approximately 75 percent (hazard ratio, 0.25;  $P = 0.03$ ). Similarly, in 30 atenolol-treated patients, the presence of diabetes was not associated with increased risk of mortality (hazard ratio,

1.2;  $P = 0.76$ ). In placebo-treated patients, the presence of diabetes was associated with a 4-fold increase in risk (hazard ratio, 4.0;  $P = 0.003$ ). No other variables were associated with outcome, including type of surgery, duration of surgery or hospitalization, and administration of  $\beta$ -  
5 blockers, calcium channel blockers or nitrates, either prior to hospital admission or following hospital discharge.

More than 85 percent of patients tolerated intravenous atenolol administration prior to and immediately following surgery, and oral administration during the postoperative period, with more than 60 percent tolerating the full dose of  
10 atenolol (10 mg intravenously or 100 mg orally) (Table 3). In approximately 10% of patients, intravenous administration of atenolol prior to or after surgery was associated with 20% or more decrease in systolic blood pressure or heart rate (Table 3); however, no patient developed systolic blood  
15 pressure  $<90$  mm Hg or heart rate  $<40$  bpm, or required therapy. Oral administration was not associated with an increased incidence of hypotension or bradycardia, or other events.

The treatment effect found in this trial cannot be attributed to inhomogeneity between groups at baseline; in fact, a larger proportion of the atenolol-treated patients  
20 had cardiovascular disease prior to surgery, and had a greater number of risk factors known to affect cardiovascular complications following surgery (Mangano et al., 1990, N. Eng. J. Med. 323:1781-1788; Goldman et al., 1977, N. Eng. J. Med. 297:845-850; Detsky et al., 1986; Arch. Intern. Med. 146:2131-2134; Hollenberg et al., 1992, JAMA 268:205-209).  
25 The results also cannot be explained by differences in surgical technique, hospitalization, or preoperative, postoperative or discharge cardiovascular medication use, specifically  $\beta$ -blockers and calcium channel blockers. A substantial portion of all variables was distributed evenly, and the variables which may not have been, such as treatments  
30 for heart failure or diabetes, were shown not to affect the conclusions of this trial.

The patient population represents approximately 10 percent of the 30 million patients undergoing noncardiac surgery (or 3 million patients), and even assuming an atenolol effect of one-fifth of the 57 percent effect found in the clinical trial disclosed herein (or 11 percent), the  
5 intensive postoperative administration of a  $\beta$ -blocker may save 33,000 lives per year at a cost of less than 100 dollars per patient (conservative estimate for one week of atenolol therapy) for the 3 million at-risk patients, or an overall cost equalling 9,000 dollars per live saved.

10 The present invention is not to be limited in scope by the exemplified embodiments, which are intended as illustrations of individual aspects of the invention. Indeed, various modifications for the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and  
15 accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

All publications cited herein are incorporated by reference in their entirety.

20

25

30

Table 1. Long-term Deaths

GROUP	PATIENT NO.	AGE	CV RISK FACTORS	TYPE OF SURGERY	TIME TO DEATH (DAYS)	CAUSE OF DEATH
5	Placebo	65	DM, HTN, PVD, AGE $\geq 65$	Peripheral Vascular	19	Massive GI hemorrhage
		67	HTN, age $\geq 65$	Peripheral Major	24	Sudden Cardiac death
		77	PVD, age $\geq 65$	AAA Repair	33	CHF, severe CAD
		64	DM, SM	Peripheral Major	35	CHF
		67	CAD, DM, HTN, PVD, age $\geq 65$	Peripheral Vascular	97	Cardiac Arrest
		65	SM, PVD, age $\geq 65$	AFBG	112	Acute bronchopneumonia, COPD
		75	DM, PVD, age $\geq 65$	Carotid	162	Sudden cardiac death
		69	HTN, age $\geq 65$	Intra-abdominal	185	Adeno CA, colon
		78	HTN, PVD, age $\geq 65$	Carotid	197	Acute MI, post PTCA
		62	CAD, HTN, PVD, SM	Peripheral Vascular	236	Acute MI
		80	CAD, HTN, age $\geq 65$	Intra Abdominal	303	Cardiac arrest
		64	CAD, SM, PVD	Peripheral Vascular	325	Sepsis
		69	CAD, DM, HTN, age $\geq 65$	Intra-abdominal	328	Small bowel obstruction 2° to CA, prostate
		77	CAD, DM, HTN, SM, PVD, age $\geq 65$	Intra-abdominal	376	Acute MI
		65	DM, SM, age $\geq 65$	Peripheral Major	384	Bladder CA
		65	CAD, DM, age $\geq 65$	Intra-abdominal	517	Sepsis 2° bowel obstruction
		75	CAD, DM, PVD, age $\geq 65$	Peripheral Vascular	517	Cardiac Arrest
		75	DM, age $\geq 65$	Intra-abdominal	629	Metastatic CA, colon
		81	DM, age $\geq 65$	Peripheral Major	658	Acute MI, ARDS
		69	DM, HTN, age $\geq 65$	Intra-abdominal	734	Post MI CVA
		66	HTN, SM, PVD, age $\geq 65$	Carotid	755	Peritonitis 2° to perforation of ileum
15	Atenolol	56	SM, PVD	Peripheral Vascular	237	Respiratory failure
		56	CAD, PVD, HTN, SM	Peripheral Vascular	295	Ventricular tachycardia
		78	CAD, DM, HTN, age $\geq 65$	Intra-abdominal	327	Severe CAD, sepsis
		66	CAD, HTN, SM, age $\geq 65$	Peripheral Major	385	Sepsis, ALS
		67	DM, THN, age $\geq 65$	Peripheral Major	416	Metastatic renal CA
		74	CAD, DM, HTN, age $\geq 65$	Peripheral Minor	481	CHF post CABG
		79	HTN, age $\geq 65$	Intra-abdominal	529	ARDS
		70	CAD, HTN, PVD, age $\geq 65$	Carotid	582	Cardiac arrest, lung CA

9 78 HTN, SM, PVD, age ≥65 Carotid 656 Metastatic squamous cell CA, layrnx, lung

CAD denotes coronary artery disease (consisting of previous CABG, MI, typical agina, chest pain with ischemic ECG responsive to exercise, scintigraphic evidence of myocardial perfusion defect, or abnormal coronary angiography), CHF congestive heart failure, COPD chronic obstructive pulmonary disease, ARDS adult respiratory distress syndrome, ALS amyotrophic lateral sclerosis, APBF aorto-femoral bypass graft, AAA abdominal aortic aneurysm, CA cancer, DM diabetes mellitus, HTN hypertension, PVD vascular disease, and SM smoking.

Table 2. Variables Associated with 30 Deaths among 200 Patients Undergoing Non-cardiac Surgery

PREDICTOR	HAZARDS		P VALUE
	RATIO	CONFIDENCE INTERVAL	
Univariable models			
Atenolol	0.4	0.2 - 0.9	0.03
Diabetes mellitus	3.1	1.4 - 6.8	0.01
Oral Hypoglycemic treatment	2.6	1.1 - 6.2	0.03
Insulin Treatment	2.6	1.0 - 6.9	0.05
Holter ischemia postop days 0-2	2.3	1.0 - 5.3	0.04
Multivariate models			
Diabetes mellitus	2.8	1.4 - 6.2	
Atenolol	0.5	0.2 - 1.1	

Table 3 - Use of Cardiovascular Medications

	PERCENT OF SUBJECTS									
	β-BLOCKER		CALCIUM CHANNEL BLOCKER		NITRATE		ACE INHIBITOR			
	ATENLOLOL	PLACEBO	ATENLOLOL	PLACEBO	ATENLOLOL	PLACEBO	ATENLOLOL	PLACEBO	ATENLOLOL	PLACEBO
Prior to Admission	18	8	22	34	8	13	22	8	22	8
		0.02 <sup>1</sup>		0.11 <sup>3</sup>		0.36		0.003 <sup>7</sup>		
5 At hospital discharge	13	7	18	27	7	15	19	6	19	6
		0.12 <sup>2</sup>		0.18 <sup>4</sup>		0.09 <sup>6</sup>		0.003 <sup>8</sup>		
At 6 months	14	8	19	30	16	22	15	18	15	18
		0.27		0.10 <sup>5</sup>		0.30		0.58		
At 12 months	17	14	24	30	19	27	24	24	24	24
		0.61		0.36		0.25		0.98		
At 24 months	16	14	19	25	14	18	18	22	18	22
		0.79		0.36		0.51		0.61		

\* - Chi-square statistics were used to test the difference between the two treatment groups.

+ - Number of atenolol patients = 99; number of placebo patients = 101.

§ - Number of atenolol patients = 94; number of placebo patients = 99.

¶ - Number of atenolol patients = 93; number of placebo patients = 91.

|| - Number of atenolol patients = 93; number of placebo patients = 91.

¥ - Number of atenolol patients = 90; number of placebo patients = 85.

1 - Odds ratio for pre-admission β-blocker use and 2-year mortality = 0.80 (P=0.73)

2 - Odds ratio for discharge β-blocker use and 2-year mortality = 0.61 (P=0.52)

3 - Odds ratio for pre-admission calcium blocker use and 2-year mortality = 1.06 (P=0.90)

4 - Odds ratio for discharge calcium blocker use and 2-year mortality = 0.85 (P=0.74)

5 - Odds ratio for 6-month calcium blocker use and 2-year mortality = 1.05 (P=0.92)

6 - Odds ratio for discharge nitrate use and 2-year mortality = 1.32 (P=0.64)

7 - Odds ratio for pre-admission ACE inhibitor use and 2-year mortality = 1.45 (P=0.50)

8 - Odds ratio for discharge ACE inhibitor use and 2-year mortality = 1.17 (P=0.79)



**WHAT IS CLAIMED IS:**

1. A method for preventing myocardial infarction in a patient following surgery, said method comprising the step of: administering to said patient a therapeutic amount of a pharmacologic cardiovascular agent after surgery, wherein said patient did not receive the pharmacologic cardiovascular agent prior to surgery for treatment of a preexisting cardiovascular condition; thereby preventing myocardial infarction in the patient.
2. The method of Claim 1 in which the agent is administered after surgery until hospital discharge.
3. The method of Claim 2 in which the agent is administered daily after surgery for at least three days.
4. The method of Claim 2 in which the agent is administered daily after surgery for up to seven days.
5. The method of Claim 1 in which the agent is a  $\beta_1$ -adrenergic selective blocking agent.
6. The method of Claim 5 in which the agent is atenolol.
7. The method of Claim 1 in which the agent is an  $\alpha$ -2 agonist.
8. The method of Claim 1 in which the agent is a nitrate.
9. The method of Claim 1 in which the agent is a calcium channel blocker.
10. The method of Claim 1 in which the agent is an ACE inhibitor.

11. The method of Claim 1 in which the agent is a platelet inhibitor.

12. The method of Claim 1 in which the agent is a thrombosis inhibitor.

5

13. The method of Claim 1 in which the surgery is cardiac-related surgery.

14. The method of Claim 1 in which the surgery is non-cardiac-related surgery.

10

15. The method of Claim 1 in which the patient suffers from coronary artery disease.

16. The method of Claim 1 in which the patient is at risk for coronary artery disease.

15

17. A method for preventing congestive heart failure in a patient following surgery, said method comprising the step of: administering to said patient a therapeutic amount of a pharmacologic cardiovascular agent after surgery, wherein said patient did not receive the pharmacologic cardiovascular agent prior to surgery for treatment of a preexisting cardiovascular condition; thereby preventing congestive heart failure in the patient.

20

18. The method of Claim 17 in which the agent is administered after surgery until hospital discharge.

25

19. The method of Claim 18 in which the agent is administered daily after surgery for at least three days.

20. The method of Claim 18 in which the agent is administered daily after surgery for up to seven days.

30

21. The method of Claim 17 in which the agent is a  $\beta_1$ -adrenergic selective blocking agent.

22. The method of Claim 21 in which the agent is atenolol.

5

23. The method of Claim 17 in which the agent is an  $\alpha$ -2 agonist.

24. The method of Claim 17 in which the agent is a nitrate.

10

25. The method of Claim 17 in which the agent is a calcium channel blocker.

26. The method of Claim 17 in which the agent is an ACE inhibitor.

15

27. The method of Claim 17 in which the agent is a platelet inhibitor.

28. The method of Claim 17 in which the agent is a thrombosis inhibitor.

20

29. The method of Claim 17 in which the surgery is cardiac-related surgery.

30. The method of Claim 17 in which the surgery is non-cardiac-related surgery.

25

31. The method of Claim 17 in which the patient suffers from coronary artery disease.

32. The method of Claim 17 in which the patient is at risk for coronary artery disease.

30

33. A method for preventing angina in a patient following surgery, said method comprising the step of: administering to said patient a therapeutic amount of a pharmacologic cardiovascular agent after surgery, wherein said patient did not receive the pharmacologic cardiovascular agent prior to surgery for treatment of a preexisting cardiovascular condition; thereby preventing angina in the patient.

34. The method of Claim 33 in which the agent is administered after surgery until hospital discharge.

35. The method of Claim 34 in which the agent is administered daily after surgery for at least three days.

36. The method of Claim 34 in which the agent is administered daily after surgery for up to seven days.

37. The method of Claim 33 in which the agent is a  $\beta_1$ -adrenergic selective blocking agent.

38. The method of Claim 37 in which the agent is atenolol.

39. The method of Claim 33 in which the agent is an  $\alpha$ -2 agonist.

40. The method of Claim 33 in which the agent is a nitrate.

41. The method of Claim 33 in which the agent is a calcium channel blocker.

42. The method of Claim 33 in which the agent is an ACE inhibitor.

43. The method of Claim 33 in which the agent is a platelet inhibitor.

44. The method of Claim 33 in which the agent is a thrombosis inhibitor.

5

45. The method of Claim 33 in which the surgery is cardiac-related surgery.

46. The method of Claim 33 in which the surgery is non-cardiac-related surgery.

10

47. The method of Claim 33 in which the patient suffers from coronary artery disease.

48. The method of Claim 33 in which the patient is at risk for coronary artery disease.--

15

20

25

30

## ABSTRACT

The present invention relates to methods for reducing mortality and cardiovascular morbidity following surgery. In particular, the invention relates to the intensive  
5 postoperative administration of a pharmacologic cardiovascular agent to reduce mortality and cardiovascular complications. The invention is illustrated by way of working examples which demonstrate that in patients with, or at risk for, coronary artery disease undergoing major noncardiac surgery, the administration of a  $\beta$ -adrenergic  
10 blocking agent throughout the period of hospitalization:  
1) reduces mortality and cardiovascular events following hospital discharge; 2) is safe and well tolerated; and 3) the estimated cost savings in lives more than outweighs the cost of therapy.

15


20

25

30


9114-003-999 (Sheet 1 of 2)

Figure 1



9114-003-999 (Sheet 2 of 2)

Figure 2





# DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below at 201 et seq. underneath my name.

I believe I am the original, first and sole inventor if only one name is listed at 201 below, or an original, first and joint inventor if plural names are listed at 201 et seq. below, of the subject matter which is claimed and for which a patent is sought on the invention entitled

## METHODS FOR REDUCING MORTALITY AND MORBIDITY BY POSTOPERATIVE ADMINISTRATION OF A PHARMACOLOGIC CARDIOVASCULAR AGENT

and for which a patent application:

- ☐ is attached hereto and includes amendment(s) filed on \_\_\_\_\_ (if applicable)  
☒ was filed in the United States on December 3, 1996 as Application No. 08/787,056 (for declaration not accompanying application)  
 with amendment(s) filed on \_\_\_\_\_ (if applicable)  
☐ was filed as PCT international Application No. \_\_\_\_\_ on \_\_\_\_\_ and was amended under PCT Article 19 on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified application, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

EARLIEST FOREIGN APPLICATION(S), IF ANY, FILED PRIOR TO THE FILING DATE OF THE APPLICATION			
APPLICATION NUMBER	COUNTRY	DATE OF FILING (day, month, year)	PRIORITY CLAIMED
			YES <input type="checkbox"/> NO <input type="checkbox"/>
			YES <input type="checkbox"/> NO <input type="checkbox"/>

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

APPLICATION NUMBER	FILING DATE

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NO.	FILING DATE	STATUS		
		PATENTED	PENDING	ABANDONED


POWER OF ATTORNEY: As a named inventor, I hereby appoint S. Leslie Misrock (Reg. No. 18872), Harry C. Jones, III (Reg. No. 20280), Berj A. Terzian (Reg. No. 20060), Gerald J. Flintoft (Reg. No. 20823), David Weild, III (Reg. No. 21094), Jonathan A. Marshall (Reg. No. 24614), Barry D. Rein (Reg. No. 22411), Stanton T. Lawrence, III (Reg. No. 25736), Isaac Jarkovsky (Reg. No. 22713), Joseph V. Colaianne (Reg. No. 20019), Charles E. McKenney (Reg. No. 22795), Philip T. Shannon (Reg. No. 24278), Francis E. Morris (Reg. No. 24615), Charles E. Miller (Reg. No. 24576), Gidon D. Stern (Reg. No. 27469), John J. Lauter, Jr. (Reg. No. 27814), Brian M. Poissant (Reg. No. 28462), Brian D. Coggio (Reg. No. 27624), Rory J. Radding (Reg. No. 28749), Stephen J. Harbulak (Reg. No. 29166), Donald J. Goodell (Reg. No. 19766), James N. Palik (Reg. No. 25510), Thomas E. Friebe (Reg. No. 29258), Laura A. Coruzzi (Reg. No. 30742), Jennifer Gordon (Reg. No. 30753), Jon R. Stark (Reg. No. 30111), Allan A. Fanucci (Reg. No. 30256), Geraldine F. Baldwin (Reg. No. 31232), Victor N. Balancia (Reg. No. 31231), Samuel B. Abrams (Reg. No. 30605), Steven I. Wallach (Reg. No. 35402), Marcia H. Sundeen (Reg. No. 30893), Paul J. Zegger (Reg. No. 33821), Edmond R. Bannon (Reg. No. 32110), Bruce J. Barker (Reg. No. 33291), Adriane M. Antler (Reg. No. 32605), Ann L. Gisolfi (Reg. No. 31956), SaraLynn Mandel (Reg. No. 31853), Mark A. Farley (Reg. No. 33170), and James G. Markey (Reg. No. 31636), all of Pennie & Edmonds LLP, whose addresses are 1155 Avenue of the Americas, New York, New York 10036, 1667 K Street N.W., Washington, DC 20006 and 3300 Hillview Avenue, Palo Alto, CA 94304, and each of them, my attorneys, to prosecute this application, and to transact all business in the Patent and Trademark Office connected therewith.

**SEND CORRESPONDENCE TO:** PENNIE & EDMONDS LLP  
1155 AVENUE OF THE AMERICAS  
NEW YORK, N.Y. 10036-2711

**DIRECT TELEPHONE CALLS TO:**  
PENNIE & EDMONDS LLP DOCKETING  
(212) 790-2803

201	FULL NAME OF INVENTOR	LAST NAME Mangano	FIRST NAME Dennis	MIDDLE NAME T.	
	RESIDENCE & CITIZENSHIP	CITY Hillsborough	STATE OR FOREIGN COUNTRY California	COUNTRY OF CITIZENSHIP U.S.A.	
	POST OFFICE ADDRESS	STREET 30 Genevra Road	CITY Hillsborough	STATE OR COUNTRY California	ZIP CODE 94010
202	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP	
	POST OFFICE ADDRESS	STREET	CITY	STATE OR COUNTRY	ZIP CODE
203	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP	
	POST OFFICE ADDRESS	STREET	CITY	STATE OR COUNTRY	ZIP CODE
204	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP	
	POST OFFICE ADDRESS	STREET	CITY	STATE OR COUNTRY	ZIP CODE
205	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP	
	POST OFFICE ADDRESS	STREET	CITY	STATE OR COUNTRY	ZIP CODE

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 201  Dennis T. Mangano	SIGNATURE OF INVENTOR 202	SIGNATURE OF INVENTOR 203
DATE 6/14/1997	DATE	DATE
SIGNATURE OF INVENTOR 204	SIGNATURE OF INVENTOR 205	SIGNATURE OF INVENTOR 206
DATE	DATE	DATE